

# Technical Data Sheet

## Eastman™ DTBHQ

### Applications

- General industrial coatings
- Intermediates
- Kitchen & bath
- Lubricants
- Molding & trim interior
- Polymer modification
- Wire/cable

### Product Description

Eastman™ DTBHQ (2,5-ditert-butylhydroquinone) is a white to tan crystalline solid that is insoluble in water and caustic solutions. Eastman™ DTBHQ has a low vapor pressure and has short induction periods in peroxide-catalyzed polymerizations.

### Typical Properties

Property	Typical Value, Units
<b>General</b>	
Molecular Weight	222.31
Empirical Formula	C <sub>14</sub> H <sub>22</sub> O <sub>2</sub>
Appearance	White to tan crystals
Assay	99 wt %
Specific Gravity	1.07
Melting Point	215 °C (419 °F)
Boiling Point	321 °C (610 °F)
Flash Point	
Cleveland Open Cup	216 °C (420 °F)
Autoignition Temperature	421 °C (790 °F)
Solubility @ 25° (77°F)	
In Acetone	25 wt %
In Caustic	Insoluble
In Ethanol (95%)	17 wt %
In Ethyl Acetate	31 wt %
In Styrene	< 5 wt %
In Toluene	< 1 wt %
in Water	Insoluble

### Comments

Properties reported here are typical of average lots. Eastman makes no representation that the material in any particular shipment will conform exactly to the values given.

*Eastman and its marketing affiliates shall not be responsible for the use of this information, or of any product, method, or apparatus mentioned, and you must make your own determination of its suitability and completeness for your own use, for the protection of the environment, and for the health and safety of your employees and purchasers of your products. No warranty is made of the merchantability of fitness of any product, and nothing*

*herein waives any of the Seller's conditions of sale.*

2/28/2018 11:35:39 AM

© 2019 Eastman Chemical Company or its subsidiaries. All rights reserved. As used herein, ® denotes registered trademark status in the U.S. only.